

TrapAll® S.W.D.

Reg. No. / Nr. L11710 Act 36 of 1947
READ THE LABEL BEFORE USE
KEEP OUT OF REACH OF CHILDREN AND ANIMALS

TrapAll® S.W.D. is a mass trapping system for the suppression of spotted wing drosophila (*Drosophila suzukii*) in berries, any other susceptible crops, home gardens and non-crop situations.



DANGER

Hazard statements

Highly Flammable liquid and vapor.
Causes severe skin burns and eye damage.

Precautionary statements

Do not breathe vapours.
Wear protective gloves.

Active ingredients
S.W.D. Volatile blend

1000 g/kg

1 Hectare kit for ____ weeks

McPhail Traps X ____

S.W.D. PheroLure® x ____

Insecticide block x ____

REGISTRATION HOLDER

Insect Science (Pty) Ltd.
Reg. Nr. 2000/022528/07
9 Industria Street, New Industrial Area, Tzaneen, 0850
Tel: +27 15 307 1391 / + 27 15 065 0199
Email: info@insectscience.co.za
www.insectscience.co.za



Batch No.:
Date of manufacture:
Expiry Date:

--

UN No. 2924

In case of poisoning, call the following number: 0861 555 777
Emergency number 082 446 8946

WARNINGS

- Handle with care.
- Keep away from children and uninformed persons and animals.
- The neat material loaded in the lures may cause severe skin burns and eye damage. When loaded into the lure, the neat material is not expected to pose significant health or environmental hazards.
- Minimize handling of lures for optimal efficacy.
- Store in a freezer and use upon opening.
- No pre-harvest interval is established for the use of **TrapAll® S.W.D.** as it is not applied directly onto the crop. No residues are expected if used according to label instructions.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions. The action and effect thereof may be affected by factors such as abnormal climatic and storage conditions such as excessive winds and rain, the occurrence of resistance of pests, and by the method and timing of application. The registration holder does not accept responsibility for damage to crops, vegetation and the environment, harm to man or animal, or for the lack of performance of the remedy concerned, due to failure of the user to follow the label instructions, or the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of uncertainty.

PRECAUTIONS

- Avoid contact with skin. Wear protective gloves.
- Take off contaminated clothing and wash it before reuse.
- IF ON SKIN: Wash with plenty of water. Get medical help if needed.
- Do not breathe vapours.
- Dispose of used lures in accordance with local regulations. Do not leave devices in the field indefinitely.

RELEVANT SUBSTANCES

S.W.D. Proprietary blend (100%)

FIRST AID

- Ingestion: The neat material is not considered to be acutely toxic by ingestion, but is considered corrosive. If swallowed, wash mouth with water if person is conscious. Never give anything by mouth to an unconscious person. If swallowed, do not induce vomiting. Obtain medical attention immediately.
- Skin: The neat material is considered to cause severe skin burns. In case of contact, remove contaminated clothing if applicable and wash area of exposure with non-abrasive soap and copious amounts of water for several minutes. Consult a physician.
- Eyes: The neat material causes severe eye damage. If product comes into contact with eyes, immediately flush eyes with clean water for several minutes, holding the eyelids open. If present, remove contact lenses after 5 minutes. Continue rinsing.
- Inhalation: The neat material is not expected to be acutely toxic by inhalation. If inhaled, remove the source of contamination or move victim to fresh air.

USE LIMITATIONS

- Under normal conditions the **TrapAll® S.W.D.** will attract *Drosophila suzukii*. for a period of approximately 4 - 6 weeks.
- The purpose of this kit is mass trapping for suppression of the spotted wing drosophila population within the field, it is not a stand-alone control measure and is ideal in an IPM programme.

DIRECTIONS FOR USE

Only use as directed.

Placement of lure and trap assembly:

- The **TrapAll® S.W.D.** is designed for use in a **McPhail Trap** or any other dry trap together with an **Insecticide block**.
- To assemble the **McPhail Trap**, push the four (4) clear inserts supplied from the outside through the four side holes in the yellow bucket until they fit tightly. These clear inserts will then point toward the inside of the yellow bucket, preventing any fruit flies entering the **McPhail Trap** to escape.
- Remove the **S.W.D. PheroLure®** and **Insecticide block** from the foil pouch and place inside the yellow bucket at the bottom. The **Insecticide block** will kill all fruit flies entering the trap.
- **DO NOT CUT OPEN THE S.W.D. PHEROLURE®.** The attractants are released through the membrane of the dispenser into the atmosphere.
- Join the clear dome lid and yellow bucket by pushing them together. Insert a trap hanger through the “ear” on the top of the clear dome lid to hang the trap with.

Placement of the trap

- Apply 12 – 120 traps per hectare, approximately 833 m² - 83 m².
- The traps should be placed evenly throughout the field or on the perimeter of the block depending on trap density.
- Suspend the **McPhail Trap** onto a trap stand or secure the trap 1.5 to 1.8 m above the ground to ensure effective attractant distribution.
- Ensure that the trap is not put in the working row to avoid mechanical damage from tractors etc.
- Remove all leaves and branches that may obstruct entrance to the trap.
- Place trap in the direction of the prevailing winds to maximise spreading of the attractant.
- Identify every trap with a number for reference and recording purposes.

Timing of trap placement

- Place traps at the beginning of the production season or as soon as pest is present.

Trap maintenance

- **S.W.D. PheroLure®** should be replaced every 4 - 6 weeks.
- Remove the used lures from the field and dispose of according to local regulations.

Monitoring method:

- It is imperative that trap catches be recorded weekly and preferably on the same day each week.
- Record the number of flies caught, the trap number and date.
- After fruit fly counts have been recorded, the caught flies must be removed from the trap.

S.W.D. PheroLure® should be used upon opening.

PheroLure® is a registered trademark of Birdberry Holdings (Pty) Ltd.

INSECT SCIENCE (Pty) Ltd.

The copyright on all documents is reserved. All unauthorized reproduction is forbidden.
